
# C21 CHEMICAL SIGNALLING



How do cells distinguish between the many different signals that they receive?

What interactions occur inside animal cells in response to chemical signals?

**Linking Questions** 

What patterns exist in communication in biological systems?

In what ways is negative feedback evident at all levels of biological organization?


2 C

Theme: Interactions + Interdependence

Level of Organization: Cells

Written and drawn by:

PETER MARIER



# HLLEARNINGOUTCOMES

C2.1.1	Receptors as proteins with binding sites for specific signalling chemicals	Students should use the term "ligand" for the signalling chemical.			
C2.1.2	Cell signalling by bacteria in quorum sensing	Include the example of bioluminescence in the marine bacterium Vibrio fischeri.			
C2.1.3	Hormones, neurotransmitters, cytokines and calcium ions as examples of functional categories of signalling chemicals in animals	Students should appreciate the differences between these categories.			
C2.1.4	Chemical diversity of hormones and neurotransmitters	Consider reasons for a wide range of chemical substances being used as signalling chemicals. Include amines, proteins and steroids as chemical groups of hormones. A range of substances can serve as neurotransmitters including amino acids, peptides, amines and nitrous oxide.			
C2.1.5	Localized and distant effects of signalling molecules	Contrasts can be drawn between hormones transported by the blood system and neurotransmitters that diffuse across a synaptic gap.			
C2.1.6	Differences between transmembrane receptors in a plasma membrane and intracellular receptors in the cytoplasm or nucleus	Include distribution of hydrophilic or hydrophobic amino acids in the receptor and whether the signalling chemical penetrates the cell or remains outside.			
C2.1.7	Initiation of signal transduction pathways by receptors	Students should understand that the binding of a signalling chemical to a receptor sets off a sequence of responses within the cell.			
C2.1.8	Transmembrane receptors for neurotransmitters and changes to membrane potential	Use the acetylcholine receptor as an example. Binding to a receptor causes the opening of an ion channel in the receptor that allows positively charged ions to diffuse into the cell. This changes the voltage across the plasma membrane, which may cause other changes.			
C2.1.9	Transmembrane receptors that activate G proteins	Students should understand how G protein-coupled receptors convey a signal into cells. They should appreciate that there are many such receptors in humans.			
C2.1.10	Mechanism of action of epinephrine (adrenaline) receptors	Include the roles of a G protein and cyclic AMP (cAMP) as the second messenger.  NOS: Students should be aware that naming conventions are an example of international cooperation in science for mutual benefit. Both "adrenaline" and "epinephrine" were coined by researchers and are based on production of the hormone by the adrenal gland; "adrenaline" comes from Latin ad = at and ren = kidney and "epinephrine" comes from old Greek epi = above and nephros = kidney, respectively. Unusually, these two terms persist in common use in different parts of the world.			
C2.1.11	Transmembrane receptors with tyrosine kinase activity	Use the protein hormone insulin as an example. Limit this to binding of insulin to a receptor in the plasma membrane, causing phosphorylation of tyrosine inside a cell. This leads to a sequence of reactions ending with movement of vesicles containing glucose transporters to the plasma membrane.			
C2.1.12	Intracellular receptors that affect gene expression	Use the steroid hormones oestradiol, progesterone and testosterone as examples. Students should understand that the signalling chemical binds to a site on a receptor, activating it. The activated receptor binds to specific DNA sequences to promote gene transcription.			
C2.1.13	Effects of the hormones oestradiol and progesterone on target cells	For oestradiol, limit to cells in the hypothalamus that secrete gonadotropin-releasing hormone. For progesterone, limit to cells in the endometrium.			
C2.1.14	Regulation of cell signalling pathways by positive and negative feedback	Limit to an understanding of the difference between these two forms of regulation and a brief outline of one example of each.			

C2.1.1 — Receptors as proteins with binding sites for specific signalling chemicals

C.2.1.2—Cell signalling by bacteria in quorum sensing C2.1.3—Hormones, neurotransmitters, cytokines and calcium ions as examples of functional categories of signalling chemicals in animals

Cells interact and communicate with each other by sending and recieving signals. The general pathway for cell-to-cell communication is as follows:













1) Sender cell releases ligand (specific signalling chemical) into extracellular space

HL

2 Ligand travels to target cell and binds to receptor (proteins with binding sites for specific ligands)

3 Binding of ligand typically induces conformational change in receptor, initiating signal transduction

(4) Sequence of responses occur in target cell

5 Ligand dissociates from binding site of receptor

X Similar to enzyme-substrate specificity, the shape of a receptor's binding site is specific (complementary) to that of a particular ligand, ensuring selective binding and activation (preventing differently-shaped ligands from binding)

Cell-to-cell communication does not only occur in multicellular organisms; Unicellular organisms (such as bacteria) can communicate and coordinate with each other within a population via chemical signalling


Quorum sensing: process of cell-to-cell communication in bacteria which enables coordinated gene expression to changing population density as a means of synchronized activity \* named 'quorum' as the process operates on a threshold-basis where a minimum amount of a chemical signal (autoinducer) is required to initiate a synchronized, group response.



Quorum sensing is performed by Vibrio fisheri, a bioluminescent bacterium species that lives mutualistically within the light organ of the Hawaiian bobtail squid: squid provides nutrients to V. fisheri and as V. fisheri population density is high within organ, they emit light, used by squid in counter-illumination camouflage in moonlight

view from below

bacterial cell density



V. fisheri release signalling

(ligand) into extracellular space


chemical - autoinducer




individuals in population, increasing


its concentration within cytoplasm











in the form of light (bioluminescence)

Signalling chemicals (ligands) in animals can be classified into four functional categories:

	Hormones	Neurotransmitters	Cytokines	Calcium ions (Ca <sup>2+</sup> )	
Type of molecule	amines, proteins, steroids	amino acids, peptides, amines, nitrous oxide	small proteins, glycoproteins	cation	
• •	endocrine cells (typically glands) secrete hormones		Many different cell types (such as those in the immune	cells pump Ca20 into extracellular space or within organelles	
	directly into the bloodstream which bind to	neurotransmitters into narrow synaptic cleft which	system) Secrete cytokines into extracellular space	(ex: sarcoplasmic reticulum), maintaining a low concentration	
Mechanism of action	receptors on the surface or within target cells	· · ·	,	within cytoplasm. Ca2+ enters cytoplasm via facilitated	
	which may be nearby or far away in the body	on postsynaptic cell (neuron, muscle, or gland)	nearby) or the same cell from which it was released	diffusion through gated ion channels, binding to receptors.	
		synaptic cleft			
	• promotes or inhibits gene expression and cell activities	• can promote or inhibit nerve impulses by generating	promotes or inhibits gene expression and alters cellular	■ can act as second messengers within cells, inducing:	
	effects long-lasting as hormones persist for	excitatory or inhibitory postsynaptic potentials	activity (ex: immune and inflammatory response)	contraction in muscle cells, neurotransmitter release by	
		effects short-lasting due to rapid removal from receptors		pre-synaptic cells and cortical and acrosome reactions	
Examples		phrine acetylcholine, GABA, dopamine, glutamate	l	● effect typically short-lasting where Ca²¹ is pumped back	

C2.1.4—Chemical diversity of hormones and neurotransmitters C2.1.5—Localized and distant effects of signalling molecules C2.1.6—Differences between transmembrane receptors in a plasma membrane and intracellular receptors in the cytoplasm or nucleus C2.1.7—Initiation of signal transduction pathways by receptors

Both hormones and neurotransmitters act as ligands in signalling pathways, yet display large diversity in their chemical composition. This may be due to signalling systems evolving repeatedly in many different organisms, resulting in a wide array of signalling chemicals, with different signalling mechanisms based on their structure and properties. Any chemical can act as a signalling chemical provided:

• their shapes and chemical properties are distinctive enough that receptors can bind with them selectively and distinguish them from other chemicals

small enough to be easily transported into and out of cells and around organism

	Types of Neurotransmitters				Types of Hormones				
	amines	amino acid	proteins /peptides	ester	gases	amines	peptides	proteins/glycoproteins	steroids
Structure	amino acid derivative	unaltered	polypeptide - chain of	organic 0	very small gaseous	amino acid derivative	short polypeptide chain	long polypeptide chain	cholesteral derivative
	(modified group)	amino acid	Several amino acids	molecule R-COR'		(modified group)	(2-50 amino acids)	(50+ amino acids)	(four fused rings)
	dopamine Justs	GABA	opioids such	acetylcholine	nitric oxide N=0:	epinephrine N	oxytocin -	insulin see	oestradiol
Examples	serotonin no	glutamate 3/	as endorphins		carbon monoxide	thyroxine	GnRH 9550-00	LH	testosterone ~
	epinephrine mo	glycine -	. 000		:C≣0:	melatonin -	ADH 6-5	FSH 7	progesterone

Depending on how far signalling molecules are transported from source to target in organisms, their effects can be localized (ex: synaptic signalling by neurotransmitters) or distant (ex: endocrine signalling by hormones)

Synaptic signalling: mode of signalling across the junction (synapse)

between two cells in the nervous system

- neurotransmitters are secreted via exocytosis from the presynaptic cell,
   diffusing across the gap (synaptic cleft), and binding to receptors on
   the postsynaptic cell membrane
- effects are very <u>localized</u> as neurotransmitters only need to diffuse a very small distance (20-40nm) and only signal the post-synaptic cell

P neural signaling C2.2

Endocrine signalling: mode of signalling where hormones travel in bloodstream to reach target

- hormones are secreted by endocrine cells into the bloodstream, traveling until they reach and bind to receptors on or in target cells
- effects may be <u>distant</u> and <u>widespread</u> as hormones
   can travel far from source to signal many different cells




Due to the hydrophobic core of plasma membranes, hydrophilic (polar, charged) ligands cannot penetrate whereas hydrophobic (non-polar) ligands can simply diffuse through readily membranes + transport B2.1

The chemical properties of ligands and their receptor (namely their hydrophobicity/hydrophobicity/ bydrophobicity/ bydrophobicity/ will dictate the type of signalling mechanism and transduction pathway - sequences of responses and interactions within the cell

#### Hydrophobic ligand signal transduction pathway

HL


- hydrophobic ligands such as steroids freely passes into target cells via simple diffusion
- once inside the cell, they bind to an intracellular receptor (either in the cytoplasm or nucleus) forming an activated receptor-ligand complex
- 3 receptor-ligand complex typically acts as a regulator of gene expression by binding to the DNA at a specific site causing transcription of a particular gene to be promoted or inhibited thus aftering cellular activity



Hydrophilic ligand signal transduction pathway

hydrophilic ligand such as amines, peptides/proteins bind to extracellular binding site on integral transmembrane receptor

- 2 binding causes conformational change in receptor structure:
- ligand-gated ion channels -> pore opens, allowing diffusion
- G protein-coupled receptors → G protein becomes activated
- receptor tyrosine Kinase -> tyrosine on tails phosphorylated
- 3 second messenger within cell synthesized/released which triggers signal transduction cascade ultimately resulting in a cellular response, such as changes in gene expression, enzyme activity (metabolic rate), and/or cell movement



Transmembrane receptor has
hydrophilic amino acids protruding
into aqueous extracellular space
(ligand binding) and cytoplasm
(cell protein interactions) and
hydrophobic amino acids in the
core (interacting with non-polar
tails) allowing membrane spanning

C2.1.8—Transmembrane receptors for neurotransmitters and changes to membrane potential C2.1.9—Transmembrane receptors that activate G proteins C2.1.10—Mechanism of action of epinephrine (adrenaline) receptors C2.1.11—Transmembrane receptors with tyrosine kinase activity

HL

(b) glycogen phosphorylase catalyzes glycogenolysis (glycogen → glucose)

\* effect of epinephrine can be amplified at each stage of the signalling cascade

as each enzyme can catalyze many reactions, thus generating multiple signals each

membranes + transport B2.1 Acetylcholine receptors: cholinergic receptors which bind the neurotransmitter Acetylcholine (ACh.). Allows the influx of positively charged ions resulting in depolarization and the propagation of an action potential in postsynaptic cell ex: at neuromuscular junctions, ACh receptors on the sarcolemma bind ACh released from motor neurons, Causing muscle fibres to cations (Na+) diffuse through channel into cell, depolarize and initiate skeletal muscle Acetylcholine binds at two binding Binding causes conformational shape change contractions @ muscle B3.3 changing voltage across plasma membrane sites on Acetylcholine receptor in receptor, opening the gate (pore) for ions G protein-coupled receptors (GPCR): transmembrane receptor that conveys signals into cells using associated Receptor Tyrosine Kinase (RTK): transmembrane receptor that conveys signals into G protein (Guanine nucleotide-binding protein) which acts as a molecular switch for signal transduction cells through enzymatic activity (Kinase), specifically by phosphorylating tyrosine from ATP • GPCR is a polypeptide with 7 membrane-spanning a helices forming 6 loops: X typically, adding phosphate (phosphorylating) a protein activates it (and vice-versa) extracellular loops for ligand binding and intracellular loops for 6 protein interaction Insulin receptor: RTK which binds protein hormone insulin in order to signal • G protein composed of 3 subunits:  $G_{\alpha}$ ,  $G_{\beta}$ ,  $G_{\gamma}$  (inactive when  $G_{\alpha}$  bound to GDP) cellular glucose uptake from blood plasma for metabolic use and storage SPCR are the largest and most diverse group of membrone receptors in eukaryotes and are involved in light, taste, and insulin released into bloodstream by pancreatic smell detection, mood regulation, immune response, nervous transmission, endocrine pathways and more! Below is the general mechanism: B cells when blood is detected to be hyperglycemic @ G protein inactive as  $G_{\infty}$  bound to GDP Ligand binds to GPCR causing conformational change in GPCR, insulin binds to insulin RTK, causing conformational change, displacing GDP from  $G_{\infty}$ , allowing GTP to bind in its place bringing the receptor's intracellular domains together into a dimer  $\bigcirc$  GTP activates G protein causing dissociation into  $G_{\alpha}$ -GTP each tyrosine Kinase catalyzes the phosphorylation of tyrosine and GBy dimer and release from receptor residues on the neighbouring receptor tail using ATP, activating it  $G_{\alpha}$ -GTP and  $G_{\beta\gamma}$  dimer activate effector proteins which relay signal via second messengers in transduction cascade 🤗 each activated phosphorylated tyrosine activates a signalling Epinephrine (Adrenaline) receptors: GPCR which bind to epinephrine (adrenaline) to stimulate a sympathetic protein, initiating an intracellular signal cascade pathway nervous response, i.e. "fight or flight" where body mobilizes energy (glucose) for skeletal muscle contractions signalling pathways ultimately result in: (1) Epinephrine binds to epinephrine receptor inducing conformational > vesicles containing glucose transporter (GLUT4) moving change which displaces GDP from G in G protein and fusing with membrane, embedding GLUTY into  $\bigcirc$  GTP binds to  $\bigcirc$  inducing dissociation into  $\bigcirc$  GTP and  $\bigcirc$  dimer into the membrane, thus increasing glucose uptake 3 G - GTP activates membrane-bound enzyme adenylyl cyclase √ promotion of glycogenesis (glucose → glycogen) adenylyl cyclase catalyzes conversion of ATP → cyclic AMP (cAMP) × inhibition of glycogenolysis (glycogen → glucose) (5) cAMP as second messenger initiates signalling cascade ultimately resulting in activation of enzyme glycogen phosphorylase

glycogen phosphorylase

NOS: Naming conventions are an example of international cooperation in science for mutual benefit. While unusual, two terms for same molecule can persist. For example:


Adrenaline = 'ad'(at) + 'ren' (Kidney) Epinephrine = 'epi' (above) + 'nephros' (Kidney) both names based on production location - adrenal gland (found above Kidney)


C2.1.12—Intracellular receptors that affect gene expression C2.1.13—Effects of the hormones oestradiol and progesterone on target cells HL C2.1.14—Regulation of cell signalling pathways by positive and negative feedback Intracellular receptors bind to hydrophobic ligands (such as steroid harmones: testosterone, destradiol, progesterone) in the cytoplasm or nucleus where they affect gene expression by binding to specific DNA sequences general signal transduction pathway for intracellular receptors: nucleus extracellular space ocytoplasm receptor-hormone complex (RHC) binds RHC regulates RNA polymerase activity such as Steroid hormone binds to site on steroid hormone (ex: testosterone) diffuses to hormone-response element (sequence promoting gene transcription, ultimately synthesizing intracellular receptor forming an receptor-hormone complex within promoter region of gene ) through the plasma membrane into cytoplasm more of a particular protein, altering cellular activity active receptor-hormone complex enters nucleus via nuclear pore Oestradiol and Progesterone are steroid female sex hormones responsible for a wide-range of effects including regulation of menstrual cycle, triggering development of female secondary sex characteristics, maintenance of reproductive tissues and \* note: oestradiol and progesterone are also naturally found in males (although in lower concentrations) and are involved in many aspects such as mood, spermatogenesis, and more. > Oestradiol plays key role in the menstrual cycle including regulation of Gonadotropin-releasing hormone (GnRH) release > Progesterane plays key role in the menstrual cycle including the development and maintanance of endometrium Oestradiol's role at the end of followlar phase of menstrual cycle Progesterone's role during luteal phase of menstrual cycle granulosa cells of developing follicle in ovary secretes Oestradiol 1) Progesterane secreted by corpus luteum in ovaries ovarian follicle 2 Oestradiol diffuses into neurons of hypothalamus and 2) Progesterane diffuses into cells of endometrium (uterine lining) Hypothalamus 3 Progesterane binds to progesterane intracellular receptor within binds to Oestradiol intracellular receptor, forming a complex (3) receptor-Oestradiol complex binds to oestradiol-response element the cytoplasm, forming a receptor-progesterone complex in DNA and promotes gene expression of signalling proteins -(4) receptor-progesterane complex moves into nucleus via nuclear pare and inducing a surge in GnRH secretion by GnRH neurons in hypothalamus binds to progesterone response element in promoter of target gene (4) GARH binds to receptors on cells in anterior pituitary and signals the (5) complex acts as transcription factor, promoting transcription of Ant. Pituitary expression and release of Luteinizing Hormone (LH) genes for cell growth and proliferation (such as growth factors) 6 endometrial cells grow and divide, causing it to thicken (5) LH promotes follicle development and Oestradiol release (positive feedback) ovulation 6 LH surge ultimately results in ovulation X if progesterone levels fall, endometrium breaks down and is shed (menstruation) Secondary oocyte Cell signalling pathways can be regulated by both negative and positive feedback where the amount of a product in a pathway can impact how much more is produced. positive feedback: end-product stimulates more of its own production -> amplifies change in system negative feedback: end-product inhibits more of its own production \_\_\_ reverses change and leads to stabilization of system @ reproduction D3.1 thermoregulation via thyroxin stimulus Ovulation via LH surge during menstrual cycle Examples: > stimulus Examples: homeostasis D3.3 blood glucose level regulation via insulin and glucagon the childbirth via oxytocin + fruit ripening via ethylene Integration of body systems C3.1 osmoregulation via ADH Integration of body systems C3.1 (+) lactation via prolactin and oxytocin heart rate and blood pressure blood pH regulation via ventilation Oestradiol and progesterone levels in menstrual cycle @ reproduction D3.1 \* typically reinforces stimulus in order to drive a process/event to completion response





#### > Textbooks

















# Simulators / Interactives





## → Articles


Bramhachari, N. P. V., & Sheela, G. M. (2018). Vibrio fischeri Symbiotically Synchronizes Bioluminescence in Marine Animals via Quorum Sensing Mechanism. In Springer eBooks (pp. 207-219). https://doi.org/10.1007/978-981-13-2429-1 13

GPCRDB: an information system for G protein-coupled receptors. Isberg, V., Vroling, B., van der Kant, R., Li, K., Vriend, G. and Gloriam, D. Nucleic Acids Research, 2014

Lights. camera. action! How the Hawaiian bobtail squid brings a creative vision to its maritime world of small big. . . (n.d.). https://www.labxchange.org/library/items/ lb:LabXchange:c4a7aa61:html:1

Pándy-Szekeres, et al. (2022). GPCRdb in 2023: state-specific structure models using AlphaFold2 and new ligand resources. Nucleic Acids Research, 51(D1), D395-D402. https://doi.org/10.1093/nar/gkac1013


### 3D models



Luciferase




Acetylcholine Receptor



Acetylcholine




**GPCR** 




G protein




Adenylyl Cyclase + G protein



Insulin receptor



receptor



Estradiol



Endocrine system