
# B2.1 Membranes and Membrane Transport



How do molecules of lipid and protein assemble into biological membranes?

What determines whether a substance can pass through a biological membrane?

#### **Linking Questions**

What processes depend on active transport in biological systems?


What are the roles of cell membranes in the interaction of a cell with its environment?

Theme: Form and Function

Level of Organization: Cells

Written and drawn by:

PETER MARIER




https://www.artstation.com/ artwork/VgyEbR

## SL Learning Content

B2.1.1	Lipid bilayers as the basis of cell membranes	Phospholipids and other amphipathic lipids naturally form continuous sheet-like bilayers in water.
B2.1.2	Lipid bilayers as barriers	Students should understand that the hydrophobic hydrocarbon chains that form the core of a membrane have low permeability to large molecules and hydrophilic particles, including ions and polar molecules, so membranes function as effective barriers between aqueous solutions.
B2.1.3	Simple diffusion across membranes	Use movement of oxygen and carbon dioxide molecules between phospholipids as an example of simple diffusion across membranes.
B2.1.4	Integral and peripheral proteins in membranes	Emphasize that membrane proteins have diverse structures, locations and functions. Integral proteins are embedded in one or both of the lipid layers of a membrane. Peripheral proteins are attached to one or other surface of the bilayer.
B2.1.5	Movement of water molecules across membranes by osmosis and the role of aquaporins	Include an explanation in terms of random movement of particles, impermeability of membranes to solutes and differences in solute concentration.
B2.1.6	Channel proteins for facilitated diffusion	Students should understand how the structure of channel proteins makes membranes selectively permeable by allowing specific ions to diffuse through when channels are open but not when they are closed.
B2.1.7	Pump proteins for active transport	Students should appreciate that pumps use energy from adenosine triphosphate (ATP) to transfer specific particles across membranes and therefore that they can move particles against a concentration gradient.
B2.1.8	Selectivity in membrane permeability	Facilitated diffusion and active transport allow selective permeability in membranes. Permeability by simple diffusion is not selective and depends only on the size and hydrophilic or hydrophobic properties of particles.
B2.1.9	Structure and function of glycoproteins and glycolipids	Limit to carbohydrate structures linked to proteins or lipids in membranes, location of carbohydrates on the extracellular side of membranes, and roles in cell adhesion and cell recognition.
B2.1.10	Fluid mosaic model of membrane structure	Students should be able to draw a two-dimensional representation of the model and include peripheral and integral proteins, glycoproteins, phospholipids and cholesterol. They should also be able to indicate hydrophobic and hydrophilic regions.

## HL Learning Content

		Unsaturated fatty acids in lipid bilayers have lower melting points, so membranes are fluid and therefore
B2.1.11	Relationships between fatty acid composition of lipid bilayers and their fluidity	flexible at temperatures experienced by a cell. Saturated fatty acids have higher melting points and make membranes stronger at higher temperatures. Students should be familiar with an example of adaptations in membrane composition in relation to habitat.
B2.1.12	Cholesterol and membrane fluidity in animal cells	Students should understand the position of cholesterol molecules in membranes and also that cholesterol acts as a modulator (adjustor) of membrane fluidity, stabilizing membranes at higher temperatures and preventing stiffening at lower temperatures.
B2.1.13	Membrane fluidity and the fusion and formation of vesicles	Include the terms "endocytosis" and "exocytosis", and examples of each process.
B2.1.14	Gated ion channels in neurons	Include nicotinic acetylcholine receptors as an example of a neurotransmitter-gated ion channel and sodium and potassium channels as examples of voltage-gated channels.
B2.1.15	Sodium-potassium pumps as an example of exchange transporters	Include the importance of these pumps in generating membrane potentials.
B2.1.16	Sodium-dependent glucose cotransporters as an example of indirect active transport	Include the importance of these cotransporters in glucose absorption by cells in the small intestine and glucose reabsorption by cells in the nephron.
B2.1.17	Adhesion of cells to form tissues	Include the term "cell-adhesion molecules" (CAMs) and the understanding that different forms of CAM are used for different types of cell–cell junction. Students are not required to have detailed knowledge of the different CAMs or junctions.




B2.1.3—Simple diffusion across membranes. B2.1.5—Movement of water molecules across membranes by osmosis and the role of aquaporins. B2.1.6—Channel proteins for facilitated diffusion. B2.1.7—Pump proteins for active transport. B2.1.8—Selectivity in membrane permeability


Many of the processes of life (such as metabolism, homeostasis, nutrition, excretion) require transport of materials in and out of cells. This can occur both passively or actively

Passive transport


· net movement of particles from an area of high concentration to low concentration without the use of ATP




- particles are constantly in random motion (due to kinetic energy) and move independent of each other but as there are more particles in areas of high concentration that can randomly move to areas of low concentration than the other way around, a net movement from high to low concentration will occur passively.
- rate can be increased by increasing concencentration gradient and membrane surface area and decreasing distance needed to move Simple diffusion passive net movement of particles from an area of high concentration to low concentration in fluids
  - small Inon-polar molecules can diffuse easily across membranes able to pass through hydrophobic core easily > able to pass in between phospholipid molecules ex: Oz , COz , steroids (testosterone and oestradiol)



- X not selective: only depends on size and water-affinity
- > Facilitated diffusion passive net movement of particles from an area of high concentration to low concentration using membrane proteins (such as channel or carrier proteins)
  - Channel proteins: integral membrane proteins which have a hydrophilic pore allowing a passage for charged (Nat, Kt, Cl, Ca2+) or polar molecules (glucose, water) to cross the membrane's hydrophobic core easily




- X channels are selectively permeable: Shape/size/charge of pore ensures specificity ex: Nat channels only allow Nat some (like voltage-gated ion channels) can be open or closed, allowing control
- Osmosis: passive net movement of water from an area of low solute concentration to high solute concentration across a semi-permeable membrane
  - membranes allow the passage of water but restrict the movement of larger or charged solutes, thus causing water to move more
  - When water is interacting with and dissolving solutes its movement is restricted be water in a dilute solution is more free and likely to move to a more concentrated solution
  - equilibrium when overall solute concentration is equal on both sides



water entered cell

Active transport: movement of particles from an area of low concentration to high concentration using ATP




- particles in an area of low concentration are less likely to move to an area of high concentration then the other way around, thus for this to occur reliably an external energy input is required
- ATP is the cell's 'energy currency' storing potential energy from cellular respiration. When hydrolyzed it releases this energy to power metabolic processes




cell respiration C1.2

- Pump proteins: transmembrane proteins which use chemical energy ATP in order to transport particles across membranes against their concentration gradient ex: sodium-potassium pump
- X Pump proteins allow selective permeability: only transport specific ions or molecules in one direction


neural signaling C2.2



Pump protein in its stable conformation is open to one side of membrane. It has high affinity to particle (.) and binds to it.




ATP is hydrolyzed to ADP, releasing energy. Energy causes a conformational change to a more unstable form, causing it to be open to the other side of the membrane



Pump is open to other side and has low affinity for particle, releasing it against its concentration gradient. Pump reverts back to its original, stable conformation - ready for reuse

Aguaporins: channel proteins which selectively allow passage of water

- greatly increases permeability to water and osmotic rate
- located in collecting ducts in Kidneys and plant roots for water absorption



Aquaporin has a very narrow pore which is positively charged only allowing water through (excluding H+)

B2.1.11—Relationships between fatty acid composition of lipid bilayers and their fluidity.

B2.1.12—Cholesterol and membrane fluidity in animal cells. B2.1.13—Membrane fluidity and the fusion and formation of vesicles. B2.1.17—Adhesion of cells to form tissues

Temperature

low 🌡

high 🌡

Temperature impacts membrane fluidity i.e. the degree of RARRR more movement less movement greater fluidity
greater permeability random movement of the components, namely phospholipids lower fluidity which can move laterally, rotate and even switch positions lower permeability Impact of phospholipid fatty acid composition on membrane fluidity > the fatty ocid composition of phospholipids can be saturated (no double bonds) or unsaturated (≥1 double bond) @ lipids BI.I Stronger pack very tightly - high density less fluid / more viscous fatty acid tails -> **T** less permeable saturated : high intermolecular forces less flexible too solid .. high melting point 100 fluid more fluid / less viscous fatty acid tails -> pack loosely - lower density unsaturated more permeable : low intermolecular forces · low melting point fluid more flexible Seneral trend: the relative percentage composition of saturated and higher percentage of higher percentage of unsaturated fats differ depending on surrounding temperature saturated fatty acids unsaturated fatty acids > Adaptation Fish living in Antarctic waters The temperate freshwater fish A. fulvescens increases (such as <u>C.hamatus</u>) have membranes its membrane unsaturated content rich in unsaturated fatty acids when lake temperatures decrease The caribou R. tarandus has different membrane fatty acid composition in different tissues, as a trend: the further away tissues are from the body core, the more unsaturated fatty acids (hooves > upper leg > body) Impact of cholesterol on membrane fluidity > cholesteral embeds itself in the plasma X cholesterol is present in animal but not plant membranes membrane as its amphipathic-OH 'head' is polar, body is non-polar fluid > cholesterol acts as a membrane fluidity modulator: Cholesteral decreases fluidity by constraining phospholipid movement - stabilizing membrane without cholesteral with cholesterol membrane Temp Cholesterol increases fluidity by disrupting solid

phospholipid tight packing - preventing stiffening

The fluidity of the membrane allows for phospholipid bilayers to change shape and fuse, and thus the formation (endocytosis) and fusion (exocytosis) of vesicles Endocytosis: the formation of a vesicle from the invagination and pinching off of a piece of the plasma membrane - bringing extracellular content within the cell fluids (pinocytosis) 

solids (phagocytosis) Process requires ATP ex: Phagocytic white blood Developing fetus takes cells and paramecia in antibodies into engulf and destroy its blood stream pathogens / food particles from the mother via the placenta using phagocytosis ACOTTO FORTING FORTING FORTING FORTING FORTING FORTING FORTING BUSINESS. vesicle Exocytosis: the fusion of vesicle a with the plasma membrane - discharging vesicle contents into the extracellular space (i.e. outside the cell). Process requires ATP ex: Neurotransmitters. Pancreas secretes Paramecia osmoregulate by (such as Acetylcholine) digestive enzymes 👺 loading and expelling excess water are released by neurons (such as lipase) using contractile vacuoles during Synaptic into the duodenum and hormones (such as insulin) into the blood transmission in order to propagate nerve impulses stream for signaling cell-adhesion molecules (CAMs) cell-surface proteins/glycoproteins which link cells with other cells forming junctions - providing structure and tissue formation > seal cells together, creating a barrier

act as a bridge, allowing communication and material exchange

connect cells to extracellular matrix, providing anchorage


B2.1.14—Gated ion channels in neurons. B2.1.15—Sodium-potassium pumps as an example of exchange HL transporters. B2.1.16—Sodium-dependent glucose cotransporters as an example of indirect active transport

Channel proteins allow for facilitated diffusion, transporting specific materials that otherwise would only diffuse slowly or not at all across a phospholipid bilayer (such as ions). While they do not control direction of flow (always a net flow high to low concentration) they can be opened or closed reversibly to control transport

Neurotransmitter-gated ion channel Transmembrane protein channels which allow the diffusion of a specific ion 'Gated' as they open or close in response to binding of a neurotransmitter

X neurotransmitters are signalling molecules which are secreted by a neuron at synapses to cause on effect ex: Acetylcholine, Glutamate, GABA, Dopamine, Serotonin, Epinephrine

Nicotinic acetylcholine receptors: gated ion channel allowing diffusion of sodium (Na+) into the cell when two acetylcholine (or nicotine) molecules bind to the receptor, causing the cell to depolarize and an action potential to be propagated



Acetylcholine is secreted from the pre-synaptic neuron into the synaptic cleft and diffuses across Before binding, receptor is closed. preventing Nat from diffusing Two Acetylcholine molecules bind to the receptor, inducing a conformational change, opening a pore and allowing Nat to diffuse through

Voltage-gated ion channel. Transmembrane protein channels which allow the diffusion of a specific ion 'Gated' as they open or close in response to a specific membrane potential (voltage)

X potential difference or voltage is the amount of electric potential energy between two points (V or JC-1) membrane potential is the difference in electric potential (V) between the inside and outside of the cell  $\rightarrow$  inside of the cell ~-70 mV compared to the outside  $\rightarrow$  generated by the sodium-potassium pump

voltage-gated sodium channel: gated ion channel allowing diffusion of sodium (Nat) into the cell when there is an increase in membrane potential (-70mV to -55 mV) causing the cell to depolarize (negative to positive)

inactivation gate Nat

activation gate closes channel, blocking Nat diffusion (-70mV) activation gate opens, allowing Not to diffuse into cell (-55mV) inactivate gate rapidly closes channel, halting Nat diffusion (30mV)

Exchange transporters: Transmembrane protein which transports different substances in opposite directions across (antiporter) the membrane against their respective gradients, thus 'exchanging' them > Sodium - Potassium pump: exchange transporter moving 3 Nat and 2 Kt against their respective (Nat/Kt-ATPase) concentration gradients across the membrane using ATP

^[Na<sup>†</sup>] ↓[K<sup>†</sup>] **↓[Na<sup>+</sup>]** ↑[K<sup>+</sup>] ↑[Na<sup>+</sup>] ↓[K<sup>+</sup>]


X as the pump undergoes unequal exchange of cations (lose 3, take in 2) it maintains a relative negative membrane potential (-70mV)

Pump binds 3 Nat from cytoplasm of cell with high affinity ATP is hydrolyzed to ADP, binding phosphate to (phosphorylating) pump Phosphorylation causes a change in pump conformation, reducing its affinity for Nat, releasing 3 Nat outside cell New pump shape binds 2 K+ from outside cell with high affinity Binding K+ triggers release of phosphate causing pump to revert to original conformation and release 2K+ into cell Pump ready for reuse.


Indirect active transport uses the energy produced by the movement of one substance down its concentration gradient to transport another against its concentration gradient. Direct ATP use not required

(symporter)


Sodium-glucose cotransporter: cotransporter protein which transports glucose against its concentration gradient into the cell using the energy from simultaneously transporting Nat down its gradient



Nat moves down its gradient binding onto cotransporter. This energy is used to bind and move glucose into cell



> voltage-gated potassium channel: gated ion channel allowing diffusion of potassium (K+) out of the cell when the membrane potential has flipped (-70 mV to 30 mV) causing the cell to repolarize (positive to negative)





activation gate closes channel, blocking Kt diffusion (-70 mV) activation gate opens, allowing K\* to diffuse out of cell (30mV) inactivate gate rapidly closes channel, halting Kt diffusion (-70 mV)





#### each resource is hyperlinked


### Textbooks

















## Simulators / Interactives










### → Articles

Benga G. (2012). The first discovered water channel protein, later called aquaporin 1: molecular characteristics, functions and medical implications. Molecular aspects of medicine, 33(5-6), 518-534. https://doi.org/10.1016/ j.mam.2012.06.001

Ernst, R., Ejsing, C. S., & Antonny, B. (2016). Homeoviscous Adaptation and the Regulation of Membrane Lipids. Journal of molecular biology, 428(24 Pt A), 4776-4791. https://doi.org/10.1016/j.jmb.2016.08.013

Janiszewska, M., Primi, M. C., & Izard, T. (2020). Cell adhesion in cancer: Beyond the migration of single cells. The Journal of biological chemistry, 295(8), 2495–2505. https://doi.org/ 10.1074/jbc.REV119.007759


## >3D models



Voltage-gated Sodium channel




Potassium channel




Sodium-Potassium Pump




Acetylcholine Receptor



Aquaporin



Glucose transporters



Phospholipid bilayer