B1.1 CARBOHYDRATES AND LIPIDS

Guiding Questions


In what ways do variations in form allow diversity of function in carbohydrates and lipids?

How do carbohydrates and lipids compare as energy storage compounds?

Linking Questions

How can compounds synthesized by living organisms accumulate and become carbon sinks?

What are the roles of oxidation and reduction in biological systems?



Theme: Form and Function

Level of Organization: Molecules

Written and drawn by:

PETER MARIER



SL LEARNING OUTCOMES


		Students should understand the nature of a covalent bond. Students should also understand that a carbon
B1.1.1	Chemical properties of a carbon atom allowing for the formation of diverse compounds upon which life is based	atom can form up to four single bonds or a combination of single and double bonds with other carbon atoms or atoms of other non-metallic elements. Include among the diversity of carbon compounds examples of molecules with branched or unbranched chains and single or multiple rings. NOS: Students should understand that scientific conventions are based on international agreement (SI
		metric unit prefixes "kilo", "centi", "milli", "micro" and "nano").
B1.1.2	Production of macromolecules by condensation reactions that link monomers to form a polymer	Students should be familiar with examples of polysaccharides, polypeptides and nucleic acids.
B1.1.3	Digestion of polymers into monomers by hydrolysis reactions	Water molecules are split to provide the -H and -OH groups that are incorporated to produce monomers, hence the name of this type of reaction.
B1.1.4	Form and function of monosaccharides	Students should be able to recognize pentoses and hexoses as monosaccharides from molecular diagrams showing them in the ring forms. Use glucose as an example of the link between the properties of a monosaccharide and how it is used, emphasizing solubility, transportability, chemical stability and the yield of energy from oxidation as properties.
B1.1.5	Polysaccharides as energy storage compounds	Include the compact nature of starch in plants and glycogen in animals due to coiling and branching during polymerization, the relative insolubility of these compounds due to large molecular size and the relative ease of adding or removing alpha-glucose monomers by condensation and hydrolysis to build or mobilize energy stores.
B1.1.6	Structure of cellulose related to its function as a structural polysaccharide in plants	Include the alternating orientation of beta-glucose monomers, giving straight chains that can be grouped in bundles and cross-linked with hydrogen bonds.
B1.1.7	Role of glycoproteins in cell–cell recognition	Include ABO antigens as an example.
B1.1.8	Hydrophobic properties of lipids	Lipids are substances in living organisms that dissolve in non-polar solvents but are only sparingly soluble in aqueous solvents. Lipids include fats, oils, waxes and steroids.
B1.1.9	Formation of triglycerides and phospholipids by condensation reactions	One glycerol molecule can link three fatty acid molecules or two fatty acid molecules and one phosphate group.
B1.1.10	Difference between saturated, monounsaturated and polyunsaturated fatty acids	Include the number of double carbon (C=C) bonds and how this affects melting point. Relate this to the prevalence of different types of fatty acids in oils and fats used for energy storage in plants and endotherms respectively.
B1.1.11	Triglycerides in adipose tissues for energy storage and thermal insulation	Students should understand that the properties of triglycerides make them suited to long-term energy storage functions. Students should be able to relate the use of triglycerides as thermal insulators to body temperature and habitat.
B1.1.12	Formation of phospholipid bilayers as a consequence of the hydrophobic and hydrophilic regions	Students should use and understand the term "amphipathic".
B1.1.13	Ability of non-polar steroids to pass through the phospholipid bilayer	Include oestradiol and testosterone as examples. Students should be able to identify compounds as steroids from molecular diagrams.

Biomolecules: molecules found in living organisms which are essential to living processes

All biomolecules contains the element carbon, this is why life is said to be "carbon-based"

what chemical properties allows carbon to form such a large variety of compounds necessary for life?

· Carbon has 4 valence electrons, allowing it to form 4 covalent bonds with other atoms




> electrostatic attraction between positive nuclei and a shared pair of negative electrons


> covalent bonds are the strongest type of bonds thus require a lot of energy to break - leading to stable compounds


Chemistry review

- Aloms are most stable when their outer (valence) shell is complete or full. They can achieve this through bonding with other atoms by either transferring (ionic) or sharing (covalent) electrons
- > The first shell can hold 2, second 8 and third 8. Carbon has 4 so readily shares these to get 8

· Carbon can share 1, 2, or 3 pairs of electrons with another atom forming:







* position of the bonds within a molecule can differ adding further variation

- · Carbon forms many molecular shapes:
- unbranched chains

Saturated fatty acid

- $H_3C-C-CH_3$
 - valine amino acid


> branched chains

> linear chains


- 2-butyne
 - unsaturated fatty acid

> bent chains

> single rings



> multi-ringed



 Carbon can form covalent bonds with other carbons and non-metal atoms, providing much diversity

>> such as Oxygen, Hydrogen, Nitrogen, Phosphorus, Sulfur

X organic molecule must contain carbon and hydrogen ex: C, H5OH (ethanol), CHy (methane) are organic COz (carbon dioxide), NH3 (ammonia) are inorganic

- · Carbon-based molecules can be short (I carbon) or very long with no real limit (some containing thousands of carbons)
- many biomolecules have functional groups attached to a carbon skeleton. These groups have different properties and react differently




based on functional groups, carbon compounds can be organised into homologous series which have similar properties

NOS: In all science disciplines unit conventions are based on international agreement -> International System of Units (51) > this allows international collaboration and facilitates replicability and comparability of experiments and data

2 types of quantities: fundamental - independent of other variables ex: m, s, kg, K, mol, A, cd ex: °C, kgm-3, ms-2, N, W, V, Pa, Hz, V derived - dependent on other variables

• When there are different groups around a carbon, they can be arranged in different, mirrored orientations, each having different biological interactions



sci notation SI prefixes centi nano milli mega symbol $10^{3-(-6)} = 10^9$

109 pm ex conversion: 0.025 km is how 0.025 Km = 25000000 pm 1 Km many micrometers? or 2.5 × 10 pm

B1.1.2—Production of macromolecules by condensation reactions that link monomers to form a polymer. B1.1.3—Digestion of polymers into monomers by hydrolysis reactions. B1.1.9—Formation of triglycerides and phospholipids by condensation reactions Biomolecules fall into 4 major groups: O carbohydrates O proteins on nucleic acids > these exist as simple single units, monomers, and can combine together to form larger more complex polymers or macromolecules the most important metabolic reactions occurring in living organisms are condensation and hydrolysis reactions X note: monomers are often not individual atoms but smaller molecules metabolism Cl.1 Condensation reaction: where two molecules are combined together forming a **●** H, O **●**-OH H-Hydrolysis reaction: where a polymer breaks down / is digested into 2 smaller, larger more complex polymer and a molecule of water less complex molecules, using a split water molecule H₂O a catabolic reaction and releases energy > named after the fact water is produced > an anabolic reaction and requires energy hydro' = water, 'lysis' = break i.e break using water Proteins @ proteins B1.2 Carbohydrates monomers: monosaccharides \ dimer: disaccharide \ polymer: polysaccharide monomers: amino acids \rightarrow dimer: dipeptide \rightarrow X: 20 different amino acids each having peptide bond alycosidic bond a different R group many ~- glucose == glycogen glucose + glucose X how many water molecules are required to form ex: ex: anabolic enzyme: ribozymes a polypeptide made of 500 amino acids ? catabolic enzymes: peptidase, pepsin, trypsin > 500-1 = 499 water molecules glucose + galactose many B-glucose === cellulose Lipids glucose + fructose alycerol + 3 fatty acids < triglyceride Nucleic acids H -C-0-C-(CH2), CH3 HO - C - (CH2), CH3 dimer: dinucleotide -> polymer: DNA / RNA monomers: nucleotides \ H-C-O-C-(CH2)0CH3 H-C-OH OH 0-P-0 > phospholipid > phosphate group + alycerol + 2 fatty acids < OH

0-P-O

nucleic acids A1.2

phosphodiester bond

catabolic enzymes: endonuclease, helicase

ex: anabolic enzymes: DNA polymerase, RNA polymerase, DNA ligase

X lipids are technically not polymers as they are not composed of repeating units but different monomers


*can all be different or the same


HO - C - (CH2) CH3


B1.1.4—Form and function of monosaccharides.


B1.1.5—Polysaccharides as energy storage compounds

B1.1.6—Structure of cellulose related to its function as a structural polysaccharide in plants
















~Textbooks












✓ Articles




3D models

alpha and beta glucose




Ribose



Deoxyribose





Maltose



Glycogen

Cellulose






Saturated and Unsaturated fatty acids





Amylopectin




Triglyceride

Amylose





Cholesterol



Glycosyltransferase

Azuma, H., Amano, T., Kamiyama, N., Takehara, N., Jingu, M., Takagi, H., Sugita, O., Kobayashi, N., Kure, T., Shimizu, T., Ishida, T., Matsumoto, M., & Sakai, H. (2022b). First-in-human phase 1 trial of hemoglobin vesicles as artificial red blood cells developed for use as a transfusion alternative. Blood Advances. 6(21), 5711-5715. https://doi.org/ 10.1182/bloodadvances.2022007977